arXiv:1212.1447v1 [cond-mat.dis-nn] 6 Dec 2012

Rare events and scaling properties in field-induced anomalous dynamics

R. Burioni,[1] G. Gradenigo,[2] A. Sarracino,[2] A. Vezzani,[3,4] and A. Vulpiani[5]

[1]*Dipartimento di Fisica e Scienza della Terra, Università degli Studi di Parma and INFN, viale G.P.Usberti 7/A, 43126 Parma, Italy*
[2]*ISC-CNR and Dipartimento di Fisica, Università Sapienza, p.le A. Moro 2, 00185 Roma, Italy*
[3] *Centro S3, CNR-Istituto di Nanoscienze, Via Campi 213A, 41125 Modena Italy*
[4]*Dipartimento di Fisica e Scienza della Terra, Università degli Studi di Parma, viale G.P.Usberti 7/A, 43126 Parma, Italy*
[5]*Dipartimento di Fisica, Università Sapienza and ISC-CNR, p.le A. Moro 2, 00185 Roma, Italy*

(Dated: December 7, 2012)

We prove that, in particular regimes of a broad class of continuous time random walks, a small external field can turn diffusion from standard into anomalous. We illustrate our findings in the trap model, a prototype of subdiffusion in disordered and glassy materials, and in the Lévy walk process, which describes superdiffusion within inhomogeneous media. For both models, in the presence of an external field, rare events induce a singular behavior in the originally Gaussian displacements distribution, giving rise to power-law tails. Remarkably, in the subdiffusive trap model, the combined effect of highly fluctuating waiting times and of a drift yields a non-Gaussian distribution characterized by long spatial tails and strong anomalous superdiffusion.

PACS numbers: 05.40.fb 02.50.Ey 05.60.-k

Large fluctuations and rare events play an important and often crucial role in many physical processes. Their contribution strongly influences physical observables in many contexts, as for instance in laser cooling [1], liposome diffusion in nematic solutions [2], heteropolymers [3], delayed chaotic systems [4] and non-equilibrium relaxation [5, 6].

In standard diffusion, governed by the Fick's law, a Gaussian probability density function (PDF), $P(x,t)$, describes the position x of particles at time t, and the fluctuation-dissipation relations provide a powerful tool to investigate fluctuations by means of response measurements [7]. In such a case, also the out-of-equilibrium response is well under control, as it is clear from the form of the PDF, which in the presence of an external field ϵ reads:

$$P_\epsilon(x,t) \simeq \frac{1}{\sqrt{2\pi D t}} \exp\left[-\frac{(x-v_\epsilon t)^2}{2Dt}\right], \qquad (1)$$

where D is the diffusion coefficient. The field induces a finite drift, $\langle x(t)\rangle_\epsilon / t \equiv v_\epsilon \neq 0$, and a *typical* displacement $x_T(t)$, defined as the maximum of the PDF, which grows as $x_T(t) = v_\epsilon t$.

Transport properties are different in the presence of large space inhomogeneities and/or several time scales, that can produce a deviation from the Gaussian diffusive behavior [5]. Large fluctuations in time can be observed in diffusion in strongly disordered systems, where particles can get trapped for a long time in a given position. For instance in this paper we will consider the trap model [5], which describes a particle diffusing with a power law distribution of waiting times,

$$p_\alpha(\tau) = \tau^{-(\alpha+1)}, \qquad (2)$$

and which is commonly studied as a prototype of slow dynamics in glassy systems [8]. Large space fluctuations characterize instead Lévy-like motion in heterogeneous and porous materials [9–13], or in turbulent flow [14], where transport is realized through increments of size l extracted from a power-law distribution $p_\alpha(l) \sim l^{-(1+\alpha)}$ [15–17].

In the absence of an external field, $\epsilon = 0$, if the tails of the steps and waiting times distributions decay slowly enough, both the trap model and the Lévy walk lead to anomalous transport, $\langle x^2(t)\rangle \sim t^{2/z}$: subdiffusive, $z > 2$, for the trap model and superdiffusive, $z < 2$, for the Lévy walk. In these conditions, motion is largely influenced by rare events coming from the algebraic tails of the $P_0(x,t)$, and, in the case where only one exponent is relevant, one can write the generalized scaling form:

$$P_0(x,t) \sim t^{-1/z} F(x/t^{1/z}). \qquad (3)$$

Here $F(y)$ is the scaling function and $\ell(t) \sim t^{1/z}$ is defined as the scaling length of the process. Assuming also a fast asymptotic decay for the tails of $F(y)$, the anomalous behavior of the mean square displacement is straightforward. However, we stress that, in general, $F(y)$ can present a slow power-law decay, and the system can feature a "strong" anomalous behavior, induced by the presence of a physical cut-off in the distribution, which gives rise to a second scaling length in the system [18].

When the fluctuations of the distributions are not too large and the tails decay sufficiently rapidly, $P_0(x,t)$ is Gaussian and standard diffusion is recovered. Moreover, the standard equilibrium fluctuation-dissipation theorem, established in general also for anomalous dynamics [19–21], holds with $\langle x(t)\rangle_\epsilon$ growing linearly in time. One could then expect to observe a regular system in all its aspects and a modified PDF analogous to (1) when an external field is applied.

In this paper, we will show that the presence of a drift can induce an anomalous non-Gaussian dynamics even for the regimes where equilibrium measurements show a standard diffusive behavior. In particular, by solving the master equations of the two processes mentioned above, we will determine the scaling forms of the PDF in the presence of an external field. As expected, we evidence that, in the region of anomalous transport, the distributions are very sensitive to the presence of a drift. More surprisingly, in the regimes where the form of the distribution is simply Gaussian at equilibrium, we find that the presence of a field may trigger large fluctuations inducing a non-Gaussian shape of $P_\epsilon(x,t)$. This happens for $1 < \alpha < 2$ in the trap model and for $2 < \alpha < 4$ in the Lévy walk: in these cases an arbitrary small external field ϵ can induce a transition from standard to *strong* anomalous superdiffusion, namely to a regime in which the PDF and its higher order momenta are characterized by different scaling lengths.

In the trap model, a particle moves with probability 1/2 form x to $x \pm \delta_0$, where, for simplicity, we take δ_0 constant. However, the main results do not change if the displacement follows a symmetric distribution with finite variance. Between two successive steps, the particle waits for a time τ extracted from Eq. (2). In the Lévy walk model again there are time intervals of duration τ extracted from the PDF in Eq. (2), but in this case the particle, during each time lag, moves at a constant velocity v_0, extracted from a symmetric distribution with finite variance, and hence performs displacements τv_0.

In the trap model, the external field is implemented by unbalancing the jump probability, i.e. setting to $(1+\epsilon)/2$ and $(1-\epsilon)/2$ the probability of jumping to the right or to the left. For the Lévy walk, in [22, 23] it has been shown that the most natural way to drive the system out of equilibrium is to apply an external field accelerating the particle during the walk, so that the distance traveled after a scattering event is $\delta = v_0\tau + \epsilon\tau^2$.

Inspired by the Gaussian case, the simplest generalization of Eq. (1) one may assume for $\epsilon \neq 0$ reads as:

$$P_\epsilon(x,t) \sim t^{-1/z} F[(x - v_\epsilon t)/t^{1/z}]. \tag{4}$$

When $\epsilon = 0$, the left-right symmetry implies that $v_\epsilon = 0$ and $F(y)$ is an even function. In standard diffusive processes, $F(y)$ is Gaussian, $z = 2$, and the drift v_ϵ vanishes linearly with ϵ. However, both in the trap model and in the Lévy walk, Eq. (4) is made slightly more complex by the occurrence of *rare events*, which change the scaling structure introducing another characteristic length. According to our results, the correct form for the PDF in the Lévy walk is:

$$P_\epsilon(x,t) \sim t^{-1/z} F[(x - v_\epsilon t)/t^{1/z}]\ \Theta\left(\epsilon t^2 - x\right), \tag{5}$$

where $\Theta(x)$ is the Heaviside step function, which accounts for the maximum displacement possible up to time t, namely one single ballistic flight. In the case of the trap model the factor $\Theta(\epsilon t^2 - x)$ in Eq. (5) has to be replaced by $\Theta(x)$, which cuts off the power law tail due to particles with large persistence time at the origin. Although these cuts do not affect the collapse of the displacement PDFs, as we will show, they are important in order to obtain the correct scaling of the moments $\langle x^n(t)\rangle_\epsilon$, where $\langle \dots \rangle_\epsilon$ denotes the average over $P_\epsilon(x,t)$.



FIG. 1: (color online) Collapse of the PDFs of the trap model, for $\alpha = 1.5$ and field $\epsilon = 0.15$ according to Eq. (4) with velocity $v_\epsilon = 0.1$. Symbols correspond to different times: $t = 10^3$ (●), $t = 2 \cdot 10^3$ (■), $t = 5 \cdot 10^3$ (◆), and $t = 10^4$ (▲). Central inset: the same data of the main figure in log-log scale, as function of $|x - v_\epsilon t|$. Notice the power law behavior $P_\epsilon(x,t) \sim x^{-(1+\alpha)}$, in agreement with the arguments reported in the text. Top left inset: collapse of the PDF for $\alpha = 1.5$ with zero drift, $\epsilon = 0$. Notice the simple Gaussian behavior, $\ell(t) \sim t^{1/2}$, in agreement with Eq. (8).

The general scaling form for $P_\epsilon(x,t)$ in the trap model and in the Lévy walk can be obtained from the master equation of the processes: this relates, at two subsequent scattering events, $P_\epsilon(x,t)$ with the function $Q_\epsilon(x,t)$, i.e. the probability of being scattered in x at time t [24]. In particular, the master equation for the trap model with v_0 extracted from a bimodal distribution, reads:

$$\begin{aligned} Q_\epsilon(x,t) &= \int \Big[\frac{1+\epsilon}{2} Q_\epsilon(x - \delta_0, t - t') \\ &+ \frac{1-\epsilon}{2} Q_\epsilon(x + \delta_0, t - t')\Big] p_\alpha(t') dt' \\ &+ \delta(t)\delta(x) \end{aligned} \tag{6}$$

with

$$P_\epsilon(x,t) = \int Q_\epsilon(x, t - t') \int_{t'}^{\infty} p_\alpha(\tau) d\tau dt', \tag{7}$$

where $\delta(x)$ is the Dirac delta. The above equation can be solved asymptotically by a Fourier transform with respect to space and time, $P(x,t) \to \tilde{P}(k,\omega)$, in the equilibrium ($\epsilon = 0$) and out-of-equilibrium ($\epsilon \neq 0$) regimes.

Without drift, $\epsilon = 0$, the results are well known [24]. In particular, in the asymptotic regime of large space and time (k and ω close to zero), for the trap model one obtains:

$$\tilde{P}_0(k,\omega) = \begin{cases} C_1\omega^{\alpha-1}/(\delta_0 k^2/2 + C_2\omega^\alpha) & \text{if} \quad 0<\alpha<1 \\ \langle t\rangle/(\delta_0 k^2/2 + i\langle t\rangle\omega) & \text{if} \quad \alpha>1, \end{cases} \tag{8}$$

where, here and in the following, C_i and $\langle t\rangle$ are constants, determined by the distribution $p_\alpha(\tau)$, and in particular $\langle t^p\rangle = \int t^p p_\alpha(t)dt$. For $\alpha > 1$ Eq. (8) corresponds to the Gaussian propagator of standard diffusion, while for $0 < \alpha < 1$ the Gaussianity is broken and the dynamics is characterized by a subdiffusive scaling length $\ell(t) \sim t^{\alpha/2}$.

Let us now evaluate the propagators of the trap model in the presence of a drift, i.e. with $\epsilon \neq 0$. For large space and time we obtain:

$$\tilde{P}_\epsilon(k,\omega) = \begin{cases} C_1\omega^{\alpha-1}/(i\epsilon\delta_0 k + C_2\omega^\alpha) & \text{if} \quad 0<\alpha<1 \\ \langle t\rangle/[i(\epsilon\delta_0 k + \langle t\rangle\omega) + C_3\omega^\alpha] & \text{if} \quad 1<\alpha<2 \\ \langle t\rangle/[\delta_0 k^2/2 + i(\epsilon\delta_0 k + \langle t\rangle\omega)] & \text{if} \quad \alpha>2. \end{cases} \tag{9}$$

Interestingly, all the regimes follow the general scaling pictures of equation (4), however several different behaviors are realized in the dynamics. For $\alpha > 2$, the standard response of a diffusive system is reproduced. In particular, $F(y)$ is a Gaussian moving at constant velocity $v_\epsilon = \epsilon\delta_0/\langle t\rangle$. For $0 < \alpha < 1$, where Gaussianity is broken already at equilibrium, $F(y)$ is clearly non-Gaussian and the characteristic length grows as $\ell(t) \sim t^\alpha$. We notice that $v_\epsilon = 0$, and hence a rigid motion of the probability with constant velocity is not observed. For $1/2 < \alpha < 1$, one finds a superdiffusive behaviour induced by the field, as discussed in [5]. In this case, due to the fast asymptotic decay of $F(y)$, the behaviour of momenta is ruled by $\ell(t)$, namely $\langle x^n(t)\rangle_\epsilon \sim \ell^n(t)$, which is an example of *weak* anomalous diffusion [18]. Comparing equations (8) and (9), for $0 < \alpha < 1$ a singular behavior at $\epsilon = 0$ is evident, i.e. $F(y)$ changes its shape, as soon as $\epsilon \neq 0$. In particular, the scaling function becomes asymmetric with respect to the origin $y = 0$, giving rise to an overall motion in the direction of the applied external field.

The most intriguing case is for $1 < \alpha < 2$, where the presence of the field turns the diffusion from standard into *strong* anomalous: now the scaling function moves at constant velocity $v_\epsilon = \epsilon\delta_0/\langle t\rangle$, however the shape of $F(y)$ is not Gaussian, and it develops a power-law tail on the left. This tail is responsible for the breaking of the relation $\langle x^n(t)\rangle_\epsilon \sim \ell^n(t)$ between momenta and scaling length of the PDF. In Fig. 1 we show the collapse of the PDFs at different times, according to the scaling (4), for the case $\alpha = 1.5$ with and without external field (see inset). Care must be taken in doing the inverse Fourier transform of $\tilde{P}_\epsilon(k,\omega)$. As suggested by numerical data and physical intuition, in real space $F(y)$ must be cut off in $y = 0$ and this provides a forbidden zone also in the plane (k,ω) in calculating $F(y)$ from $\tilde{P}_\epsilon(k,\omega)$. Equation (9) implies then that the scaling length of the moving scaling function grows in a superdiffusive way, and in particular $z = \alpha < 2$. The superdiffusive spreading of the PDF that one can see in Fig. 1 is triggered by the rare events corresponding to particles that at the observational time t are still at the origin. One can easily understand that the persistence of particles in the same position is directly related to the waiting time PDF (2). From this observation follows that $P_\epsilon(x,t) \sim t|x - v_\epsilon t|^{-(1+\alpha)}\Theta(x)$ (see central inset of Fig. 1). The cut $\Theta(x)$ in the power law tail is necessary because at large times, even if particles have a non-negligible probability of staying close to the origin, negative values of x are almost forbidden, due to the positive field. By considering the joint effect of the power-law behavior of the left tail and of the cut-off in zero, one can obtain the spreading around the peak of the PDF: $\langle[\delta x(t)]^2\rangle_\epsilon = \langle x^2(t)\rangle_\epsilon - \langle x(t)\rangle_\epsilon^2 \sim t^{3-\alpha}$. The strong anomalous behaviour is then clear because one has $\langle[\delta x(t)]^2\rangle_\epsilon \neq \ell^2(t) \sim t^{2/\alpha}$.

Let us now consider the Lévy walk. In this case the master equation is:

$$\begin{aligned} Q_\epsilon(x,t) \;=\; & \int \Big[\frac{1}{2}Q_\epsilon(x - v_0t' - \epsilon t'^2, t-t') \\ & + \frac{1}{2}Q_\epsilon(x + v_0t' - \epsilon t'^2, t-t')\Big]p_\alpha(t')dt' \\ & + \delta(t)\delta(x) \end{aligned} \tag{10}$$

with

$$\begin{aligned} P_\epsilon(x,t) \;=\; & \int \Big[\frac{1}{2}Q_\epsilon(x - v_0t' - \epsilon t'^2, t-t') \\ & + \frac{1}{2}Q_\epsilon(x + v_0t' - \epsilon t'^2, t-t')\Big] \int_{t'}^{\infty} p_\alpha(\tau)d\tau dt'. \end{aligned} \tag{11}$$

At equilibrium, the Fourier transform $\tilde{P}_0(k,\omega)$ reads:

$$\tilde{P}_0(k,\omega) = \begin{cases} C_4\omega^{\alpha-1}/(C_5 k^\alpha + C_6\omega^\alpha) & \text{if} \quad 0<\alpha<1 \\ \langle t\rangle/(C_7 k^\alpha + i\langle t\rangle\omega) & \text{if} \quad 1<\alpha<2 \\ \langle t\rangle/(v_0^2\langle t^2\rangle k^2/2 + i\langle t\rangle\omega) & \text{if} \quad \alpha>2. \end{cases} \tag{12}$$

In this case, standard Gaussian diffusion is recovered for $\alpha > 2$, while for $1 < \alpha < 2$ and $0 < \alpha < 1$ the non-Gaussian propagator corresponds to a superdiffusive and a ballistic motion, i.e. the scaling length grows as $t^{1/\alpha}$ and as t, respectively.

In the presence of a drift, from Eqs. (10) and (11), in the asymptotic regimes where $v_\epsilon t \ll t^{2/\alpha}$ we obtain for the Fourier transform $\tilde{P}_\epsilon(k,\omega)$:

$$\tilde{P}_\epsilon(k,\omega) = \begin{cases} C_8\omega^{\alpha-1}/(C_9 k^{\alpha/2} + C_{10}\omega^\alpha) & \text{if} \quad 0<\alpha<1 \\ \langle t\rangle/(C_9 k^{\alpha/2} + i\langle t\rangle\omega) & \text{if} \quad 1<\alpha<2 \\ \langle t\rangle/[C_9 k^{\alpha/2} + i(\epsilon\langle t^2\rangle k + \langle t\rangle\omega)] & \text{if } 2<\alpha<4 \\ \langle t\rangle/[\langle t^2\rangle v_0^2 k^2/2 + i(\epsilon\langle t^2\rangle k + \langle t\rangle\omega)] & \text{if } \alpha>4 \end{cases} \tag{13}$$

From the above expressions one obtains the PDF in space and time by inverse Fourier transform, taking care also in this case of a forbidden region in the plane (k, ω). Indeed, at each observational time t, the largest displacement possible is $x = v\tau + \epsilon\tau^2 \sim \epsilon\tau^2$, corresponding to the rare event of a particle never colliding up to $\tau = t$, and this bound necessarily requires for the $\Theta(\epsilon t^2 - x)$ in Eq. (5). Correspondingly the integration in the conjugated space must be taken with the constraint $|k| \geq \omega^2/\epsilon$. With this *caveat* we find that for $\alpha > 4$ Eq. (13) describes a Gaussian distribution, moving at constant velocity $v_\epsilon = \epsilon\langle t^2\rangle/\langle t\rangle$. For $\alpha < 2$, the rigid translation of the scaling function is subdominant ($v_\epsilon t \ll t^{2/\alpha}$) and the global effect of the field is a strong asymmetry of $F(y)$ (see Fig. 2). In this case the scaling length grows in super-ballistic way, i.e. $z = 1/2$ (linearly accelerated motion) for $0 < \alpha < 1$ and $z = \alpha/2$ for $1 < \alpha < 2$. Finally, for $2 < \alpha < 4$ the drift with constant velocity v_ϵ becomes relevant, but the scaling function is still not Gaussian due to an emerging power-law tail which produces a strong anomalous superdiffusive behavior. The remarkable feature we discovered for the Lévy walk with $2 < \alpha < 4$ is that, similarly to what happens for the trap model with $1 < \alpha < 2$, the field of intensity ϵ produces a power-law tail otherwise not present in the zero-field Gaussian distribution (see Fig. 3). Moreover the scaling lenght $\ell(t)$ which yields the collapse of the PDFs does not govern the behaviour of momenta, $\langle x^n(t)\rangle_\epsilon \neq \ell^n(t)$. One can obtain that the probability of the largest displacement x at time t goes like $\sim x^{-(1+\alpha/2)}$ and such a power-law behavior of the tail can be clearly seen in Figs. 2 and 3. Indeed this power-law behavior, together with the cut-off $\Theta(\epsilon t^2 - x)$, provides for the spreading around the peak of the distribution: $\langle[\delta x(t)]^2\rangle_\epsilon = \langle x^2(t)\rangle_\epsilon - \langle x(t)\rangle_\epsilon^2 \sim t^{5-\alpha} \neq \ell(t)^2 \sim t^{4/\alpha}$, showing what claimed above: a single scaling length cannot capture the behavior of all momenta.

In this letter we have presented results for the trap model and the Lévy walk in the presence of currents. At variance with standard Gaussian systems, in such models the external field, even arbitrarily small, significantly modifies the scaling properties of the PDF of displacements, introducing a new lengthscale related to rare events. Beyond the principal scaling length of the distribution $\ell(t)$ and the one related to the rigid shift (if present), one must also consider the typical length introduced by the cut-off of the power-law tail, necessary for the calculation of higher order momenta. That is how rare events induce the observed *strong* anomalous behaviour.

In conclusion we have shown that the modification of transport properties in the presence of an external field represents a valuable probe to unveil the underlying dynamical structure of the system. In particular, field-induced anomalous behaviors highlight the importance of rare and large fluctuations in regimes where the diffusional properties are *apparently* standard.



FIG. 2: (color online) Collapse of PDFs with the scaling $t^{2/\alpha}P_\epsilon(x/t^{2/\alpha})$, for the Lévy walk with exponent $\alpha = 1.5$ and field $\epsilon = 0.25$. Inset: $P_\epsilon(x,t)$ at different times. Notice that the approximation $(x - v_\epsilon t)/t^{2/\alpha} \sim x/t^{2/\alpha}$ works very well asymptotically.



FIG. 3: (color online) Collapse of the PDFs according to the scaling $t^{2/\alpha}P_\epsilon[(x - v_\epsilon t)/t^{2/\alpha}]$, for the Lévy walk with exponent $\alpha = 2.5$ and field $\epsilon = 0.25$, with $v_\epsilon = 0.38$. Symbols correspond to different times: $t = 1.6 \cdot 10^4$ (●), $t = 3.2 \cdot 10^4$ (■), $t = 6.4 \cdot 10^4$ (♦), and $t = 1.28 \cdot 10^5$ (▲). Inset: the same data of the main figure are reported in log-log scale to highlight the power-law tail of the distribution.

We thank A. Puglisi and D. Villamaina for useful discussions. The work of GG and AS is supported by the Granular Chaos project, funded by the Italian MIUR under the grant number RIBD08Z9JE.

[1] F. Bardou, J.-P. Bouchaud, A. Aspect, and C. Cohen-Tannoudji, *Lévy Statistics and Laser Cooling: How*

Rare Events Bring Atoms to Rest (Cambridge University Press, Cambridge, 2001).

[2] B. Wang, J. Kuo, S. C. Bae, and S. Granick, Nature Materials **11**, 485 (2012).

[3] L. H. Tang and H. Chaté, Phys. Rev. Lett. **86**, 830 (2001).

[4] A. D. Sánchez, J. M. López, M. A. Rodríguez, and M. A. Matías, Phys. Rev. Lett. **92**, 204101 (2004).

[5] J.-P. Bouchaud and A. Georges, Phys. Rep. **195** 127 (1990).

[6] P. Ribière, P. Richard, R. Delannay, D. Bideau, M. Toiya, and W. Losert, Phys. Rev. Lett. **95**, 268001 (2005).

[7] U. Marini Bettolo Marconi, A. Puglisi, L. Rondoni, and A. Vulpiani, Phys. Rep. **461**, 111 (2008).

[8] L. Angelani, R. Di Leonardo, G. Parisi, and G. Ruocco, Phys. Rev. Lett. **87**, 055502 (2001).

[9] P. Levitz, Europhys. Lett. **39**, 593 (1997).

[10] D. ben-Avraham and S. Havlin, *Diffusion and Reactions in Fractals and Disordered Systems* (Cambridge University Press, Cambridge, 2004).

[11] R. Burioni, L. Caniparoli, and A. Vezzani, Phys. Rev. E **81**, 060101 (2010).

[12] P. Buonsante, R. Burioni and A. Vezzani, Phys. Rev. E **84**, 021105 (2011).

[13] O. Bénichou, C. Loverdo, M. Moreau, and R. Voituriez, Rev. Mod. Phys. **83**, 81 (2011)

[14] M. F. Shlesinger and J. Klafter, Phys. Rev. Lett., **54** 2551 (1985).

[15] P. Barthelemy, J. Bertolotti and D. S. Wiersma, Nature **453** 495 (2008).

[16] J. Bertolotti, K. Vynck, L. Pattelli, P. Barthelemy, S. Lepri, and D. S. Wiersma, Adv. Funct. Mat. **20** 6, 965 (2010).

[17] R. Klages, G. Radons and I. M. Sokolov (Eds.) *Anomalous Transport: Foundations and Applications* (Wiley, VCH Berlin, 2008).

[18] P. Castiglione, A. Mazzino, P. Muratore-Ginanneschi, and A. Vulpiani, Physica D: Nonlinear Phenomena **134**, 75 (1999).

[19] E. Barkai and V. Fleurov, Phys. Rev. E **58** 1296 (1998)

[20] S. Jespersen, R. Metzler, and H. C. Fogedby, Phys. Rev. E **59**, 2736 (1999)

[21] D. Villamaina, A. Sarracino, G. Gradenigo, A. Puglisi and A. Vulpiani, J. Stat. Mech. (2011) L01002

[22] I. M. Sokolov and R. Metzler, Phys. Rev. E **67**, 010101(R) (2003).

[23] G. Gradenigo, A. Sarracino, D. Villamaina, and A. Vulpiani, J. Stat. Mech. (2012) L06001.

[24] J. Klafter and I. M. Sokolov, *First steps in random walks* (Oxford University Press, Oxford, 2011).